

The Geological Society of America  
Field Guide 15  
2009

## ***Mesozoic sedimentation, magmatism, and tectonics in the Blue Mountains Province, northeastern Oregon***

**Todd A. LaMaskin**

*Department of Geological Sciences, University of Oregon, Eugene, Oregon 97405, USA*

**Joshua J. Schwartz**

*Department of Geological Sciences, University of Alabama, Box 870338 Tuscaloosa, Alabama 35487, USA*

**Rebecca J. Dorsey**

*Department of Geological Sciences, University of Oregon, Eugene, Oregon 97405, USA*

**Arthur W. Snoke**

*Department of Geology and Geophysics, University of Wyoming, Laramie, Wyoming 82071, USA*

**Kenneth Johnson**

*Department of Natural Sciences, University of Houston–Downtown, Houston, Texas 77002, USA*

**Jeffery D. Vervoort**

*School of Earth and Environmental Sciences, Washington State University, Pullman, Washington 99164, USA*

### **ABSTRACT**

This field trip guide describes a two-day excursion through Mesozoic accreted terranes of the Blue Mountains Province in northeastern Oregon. Day 1 is focused on sedimentary rocks of the Izee terrane. These deposits are divided into two unconformity-bounded megasequences, MS-1 and MS-2, that record two stages of syntectonic basin formation. MS-1 (Late Triassic to Early Jurassic) accumulated in fault-bounded marine sub-basins on the flank of an inferred growing Baker terrane thrust belt. MS-1 sandstones, derived from the Baker terrane, contain abundant Paleozoic, Late Paleoproterozoic, and Late Archean detrital-zircon grains. These observations suggest affinity of the Baker terrane and MS-1 in the Izee area to portions of the Klamath and Sierra Nevada terranes that contain similar detrital-zircon age distributions. MS-2 (Early to early-Late Jurassic) accumulated in a large marine basin that received input from low-grade metavolcanic rocks to the east (modern coordinates). Detrital zircons are dominated by Mesozoic, Neoproterozoic, and Mesoproterozoic grains. Two possible interpretations for MS-2 are: (1) the Jurassic Izee basin was fed directly by the large Mesozoic trans-cratonal sediment-dispersal system, or (2) trans-cratonal sediment was deposited in a Triassic backarc basin in Nevada and was later recycled into the Jurassic Izee basin during Cordilleran orogenesis.

LaMaskin, T.A., Schwartz, J.J., Dorsey, R.J., Snoke, A.W., Johnson, K., and Vervoort, J.D., 2009, Mesozoic sedimentation, magmatism, and tectonics in the Blue Mountains Province, northeastern Oregon, in O'Connor, J.E., Dorsey, R.J., and Madin, I.P., eds., *Volcanoes to Vineyards: Geologic Field Trips through the Dynamic Landscape of the Pacific Northwest: Geological Society of America Field Guide 15*, p. 187–202, doi: 10.1130/2009.fld015(09). For permission to copy, contact editing@geosociety.org. ©2009 The Geological Society of America. All rights reserved.

**Day 2 of the field trip is focused on Jurassic–Cretaceous magmatism in the Baker terrane. Late Middle Jurassic to Early Cretaceous igneous rocks in the Blue Mountains Province record three distinct pulses of plutonism that are characterized by distinctive spatial and geochemical signatures. These episodes consist of: (1) late Middle to Late Jurassic small gabbro to quartz diorite plutons (ca. 162–154 Ma; low Sr/Y); (2) Late Jurassic to Early Cretaceous plutons and batholiths (ca. 148 and 137 Ma; includes spatially distinct belts of low and high Sr/Y at 147–145 Ma); and (3) Early Cretaceous small plutons of tonalitic and trondhjemitic composition (ca. 124–111 Ma). Temporal transitions in geochemical characteristics between these suites raise fundamental questions regarding the origins of plutonism in the Baker terrane. In particular, the transition from low Sr/Y (group 1) to high Sr/Y (group 2) magmatism in the Greenhorn subterrane occurred ~ 7 Ma after regional contraction, and may record partial melting of thickened crust as a direct result of Late Jurassic orogenesis.**

## INTRODUCTION

This field trip examines Mesozoic rocks in northeastern Oregon that make up part of the Blue Mountains Province, an allochthonous group of variably metamorphosed, arc- and non-arc-related rocks in Oregon, Idaho, and Washington (Fig. 1) (Hamilton, 1963; Armstrong et al., 1977; Vallier, 1977; 1995; Dickinson and Thayer, 1978; Brooks and Vallier, 1978; Silberling et al., 1984). Recent stratigraphic analysis (Dorsey and LaMaskin, 2007, 2008) and new geochronologic and geochemical data (Johnson et al., 1995; Johnson and Barnes, 2002; LaMaskin et al., 2008; Parker et al., 2008; Schwartz and Snoko, 2008; Unruh et al., 2008; LaMaskin, 2009; Schwartz et al., 2010) challenge traditional models and suggest intriguing new ideas for tectonic and magmatic development of the Blue Mountains region. The primary objectives of this field trip are to familiarize participants with fundamental tectonic problems in the region, explore the implications of new data that are being generated in these studies, and assess how the Blue Mountains Province may fit into the larger context of Cordilleran tectonics, including relationships with southern British Columbia terranes to the north, Klamath-Sierran terranes to the south, and a Mesozoic thrust belt in western Nevada. Day 1 of the trip will focus on the boundary zone between the Baker terrane and younger sedimentary rocks of the Triassic–Jurassic Izee terrane. Day 2 of the trip will focus on Jurassic magmatism and related rocks within the Baker terrane. An important goal of the trip is to integrate information and emerging new data from the two study areas.

Rocks of the Blue Mountains Province are commonly divided into two late Paleozoic to early Mesozoic volcanic island-arc assemblages (the Wallowa and Olds Ferry terranes), a Paleozoic to early Mesozoic subduction-accretionary complex (Baker terrane), and a Triassic–Jurassic clastic sedimentary succession (Izee terrane). The *Wallowa terrane* (Fig. 1) is a succession of island-arc related plutonic, volcanic, and sedimentary rocks of Permian through Early Jurassic age (Vallier, 1977, 1995; Walker, 1986, 1995). The *Olds Ferry terrane* (Fig. 1) contains island-arc related Upper Triassic to Lower Jurassic, volcanic, and marine volcanoclastic and epiclastic rocks, as well as small Middle (ca.

235 Ma) and Upper Triassic (ca. 218–212 Ma) mafic to felsic intrusions (Brooks and Vallier, 1978; Vallier, 1995; Walker, 1986, 1995; Tumpane et al., 2008; LaMaskin, 2008; Unruh et al., 2008). The *Baker terrane* accretionary-subduction complex (Fig. 1) is situated between the Wallowa and Olds Ferry terranes and includes both island-arc and non-arc-related rocks (Jones et al., 1976; Hotz et al., 1977; Carpenter and Walker, 1992; Ferns and Brooks, 1995; Leeman et al., 1995; Vallier, 1995; Schwartz et al., 2010). At least four subterrane-level units within the Baker terrane include the Grindstone, Bourne, and Greenhorn subterrane, and the Burnt River Schist (Kays et al., 1987; Ashley, 1995; Ferns and Brooks, 1995; Schwartz et al., 2006; Schwartz and Snoko, 2008). Sedimentary units in the Baker terrane include Devonian through Triassic clastic and carbonate successions with minor occurrences of Jurassic strata. Rocks traditionally included in the *Izee terrane* (Fig. 1) consist of Triassic and Jurassic dominantly sedimentary rocks that are in depositional or fault contact with rocks of the Baker terrane in central Oregon. Correlation of Middle Jurassic rocks of the Izee terrane with the Coon Hollow Formation in the Wallowa terrane suggests that the Izee terrane represents a stratigraphic overlap assemblage that links the Blue Mountains terranes





Figure 1. Geologic map of the Blue Mountains Province, modified from LaMaskin (2009). Ages of plutons shown where data are available; z—U-Pb zircon age; K-Ar—potassium-argon age. Question marks indicate uncertain terrane affiliations and/or terrane boundary locations. The Bourne and Greenhorn subterrane, Grindstone terrane, and Burnt River Schist are here considered sub-terrane level units of the Baker terrane. MS—megasequence (after Dorsey and LaMaskin, 2007); BRS—Burnt River Schist; BMB—Bald Mountain batholith; WB—Wallowa batholith; PCF—Poison Creek fault; Cpx.—complex; Ldg.—landing; WA—Washington; OR—Oregon; ID—Idaho. Compiled from numerous sources including Dickinson and Vigrass (1965); Brown and Thayer (1966; 1977); Thayer and Brown (1966); Hendrickson et al. (1972); Brooks et al. (1976); Dickinson and Thayer (1978); Brooks (1979); Walker and MacLeod (1991); Walker (1986; 1995); Vallier (1995; 1998); Ashley (1995); Ferns and Brooks (1995); Leeman et al. (1995); Ferns et al., (2001); Lewis (2002); Lund (2004); Kays et al. (2006); Dorsey and LaMaskin (2007); Mann and Vallier (2007); Parker et al. (2008); Unruh et al. (2008); J. Schwartz (unpublished data); K. Johnson (unpublished data).



by Early to Middle Jurassic time (i.e., Pessagno and Blome, 1986; White et al., 1992; Dorsey and LaMaskin, 2007, 2008; LaMaskin et al., 2008). Regional contractile deformation affected all rocks of the Blue Mountains Province in Late Jurassic time (Avé Lallemant, 1995), and terrane amalgamation prior to ca. 144 Ma is suggested by Late Jurassic to Early Cretaceous granodioritic stitching plutons (Fig. 1; Walker, 1986, 1989).

Historically, some workers have interpreted rocks of the Blue Mountains Province to represent a single, complex far-traveled island arc (e.g., Pessagno and Blome, 1986; White et al., 1992; Vallier, 1995). Others have suggested that the Blue Mountains Province contains both intraoceanic (Wallowa) and continent-fringing (Olds Ferry) island-arc systems that are separated by a subduction/accretionary complex (Baker terrane) (e.g., Dickinson, 1979, 2004). Recent work by Dorsey and LaMaskin (2007), LaMaskin et al. (2008), and Schwartz et al. (2010) suggests that this region underwent an early Mesozoic, Molucca-Sea style arc-arc collision that involved portions of the Wallowa, Baker, and Olds Ferry terranes. In addition, Dorsey and LaMaskin (2007, 2008) and LaMaskin et al. (2008) proposed that Jurassic sedimentary rocks across the region were deposited in a subsiding flexural basin adjacent to a growing Jurassic orogenic belt in western Nevada (Luning-Fencemaker fold-and-thrust belt). The diversity of tectonic models reflects the geologic complexity in the region and ongoing attempts to interpret emerging new data using comparisons to present-day plate-tectonic settings.

## STRATIGRAPHY AND STRUCTURE OF THE IZEE AREA

Paleozoic–Mesozoic serpentinite-matrix mélange of the Baker terrane and Triassic to Jurassic sedimentary rocks of the Izee terrane are exposed in a large erosional inlier southwest of the town of John Day, Oregon (Figs. 1 and 2A; Dickinson and Vigrass, 1965; Dickinson and Thayer, 1978). Triassic and Jurassic sedimentary rocks can be divided into two regional-scale, unconformity-bounded megasequences (Dorsey and LaMaskin, 2007). Rocks of megasequence 1 include Upper Triassic to Lower Jurassic marine strata that experienced significant syndepositional deformation and are derived dominantly from the Baker terrane (i.e., an outboard provenance; Dickinson and Thayer, 1978; Dorsey and LaMaskin, 2007). Deposits of megasequence 1 occur in two distinct packages located east and west of the southeast-directed Poison Creek reverse fault (Figs. 2A and 2B). Units west of the Poison Creek fault include the Upper Triassic Vester Formation (~4000 m) and small erosional remnants of the overlying Late Triassic Rail Cabin Argillite (~600 m) and Early Jurassic Graylock Formation (~150 m). These marine deposits rest nonconformably on, and are locally infolded and faulted with, serpentinite-matrix mélange of the Baker terrane (Grindstone and Greenhorn subterrane; Dickinson and Vigrass, 1965; Dickinson and Thayer, 1978; Ferns and Brooks, 1995). East of the Poison Creek fault, sedimentary rocks of megasequence 1 are age-equivalent to the Rail Cabin Argillite and Graylock Forma-

tion and include a thick package of marine argillite and turbidites called the Aldrich Mountains Group (~10,000 m; Dickinson and Thayer, 1978). All rocks of megasequence 1 were affected by Late Triassic to Early Jurassic deformation and are overlain by rocks of megasequence 2 along a regional angular unconformity.

Sedimentary rocks of megasequence 2 in the Izee area include the Lower to Upper(?) Jurassic Mowich Group (~500 m) and Snowshoe (~1000 m), Trowbridge (~1000 m) and Lonesome formations (~3000 m; Fig. 2B). These deposits accumulated in a marine basin that experienced regional transgression from Early through Middle Jurassic time (ca. 190–161 Ma; Mowich Group through Trowbridge Formation) followed by regional regression in Middle and Late(?) Jurassic time (Lonesome Formation; Dickinson 1979; Dickinson et al., 1979). All deposits of megasequences 1 and 2 were affected by Late Jurassic contractile deformation (Avé Lallemant, 1995).

## DETRITAL-ZIRCON DATA

Detrital-zircon data from rocks of megasequence 1 and megasequence 2 in the Izee area record a major shift in provenance from Late Triassic to Middle Jurassic time, consistent with provenance trends in sandstone detrital modes identified by Dickinson et al. (1979). Late Triassic sediments of the Vester Formation are interpreted to be derived from the Baker terrane (Dickinson and Thayer, 1978), and the detrital-zircon age distributions are dominated by Paleozoic, Late Paleoproterozoic, and Late Archean ages (LaMaskin, 2009). Crystalline basement rocks older than Permian are not known to be present in the Baker terrane, and thus detrital-zircon grains in megasequence 1 were likely reworked from Paleozoic and Mesozoic(?) clastic rocks of the Baker terrane. In contrast, the Jurassic deposits are inferred to be derived from inboard sources (Dickinson and Thayer, 1978). Detrital-zircon age distributions in Jurassic samples are dominated by Mesozoic ages and include significant quantities of Neoproterozoic and Mesoproterozoic ages with lesser amounts of Late Paleoproterozoic ages.

Other accretionary-subduction complexes within terranes of the Klamath Mountains and Sierra Nevada are dominated by Paleozoic, Late Paleoproterozoic, and Late Archean detrital-zircon grains such as those eroded from the Baker terrane and deposited as megasequence 1 in the Izee region (e.g., Harding et al., 2000; Scherer, 2006). This similarity in detrital-zircon age distributions suggests that pre-Triassic portions of the Baker terrane and megasequence-1 sediments have a genetic affinity to portions of the Klamath and Sierra Nevada terranes.

In Jurassic samples, significant quantities of Neoproterozoic and Mesoproterozoic detrital-zircon grains with lesser amounts of Late Paleoproterozoic grains indicate derivation from a previously documented trans-cratonal sediment-dispersal system during Jurassic time (i.e., Rahl et al., 2003; Dickinson and Gehrels, 2003, 2009), and suggest proximity of the Jurassic Izee basin to the Cordilleran margin. Alternatively, these detrital-zircon grains may have originally been deposited in the Triassic







Figure 2. (A) Geologic map of the Izee area, modified from Dickinson and Thayer (1978), showing location of stops for Day 1. (B) Stratigraphic reconstruction of Izee area prior to Late Jurassic deformation, projected into E-W line of section (in A) using maps and data from Dickinson and Thayer (1978), Dickinson and Viggras (1965), and Brown and Thayer (1966). Total structural relief of east- to southeast-directed thrust belt is ~10–15 km based on thickness of Triassic–Jurassic syn- to post-thrusting stratigraphic section. CCF—Camp Creek fault; PCF—Poison Creek fault. Black zircon crystals and sample numbers indicate approximate position of detrital-zircon samples discussed in text.

back-arc basin in Nevada (i.e., Dickinson and Gehrels, 2008) and subsequently eroded from the Luning-Fencemaker fold-and-thrust belt during Jurassic uplift and erosion (Wyld, 2002; Wyld et al., 1996, 2003). Both explanations support a model for moderate post-Jurassic northward translation (~400 km) of the Blue Mountains Province (e.g., Wyld and Wright, 2001).

**JURASSIC–EARLY CRETACEOUS MAGMATISM IN THE BAKER TERRANE**

Late Middle Jurassic to Early Cretaceous magmatism in the Blue Mountains Province occurred in three distinct pulses of plutonism between 162 and 154 Ma, 148 and 137 Ma, and 124 and 111 Ma (Walker, 1986, 1989, 1995; Johnson et al., 1995; Johnson and Barnes, 2002; Parker et al., 2008; Schwartz and Snoke, 2008; Unruh et al., 2008; J. Schwartz and K. Johnson, unpublished data; Fig. 3; Table 1). These plutons define spatially distinct belts that parallel major tectonic features of the amalgamated Blue Mountains terranes (Wallowa, Baker, and Izee terranes), and formed prior to ~60° of post-Cretaceous clockwise rotation (Wilson and Cox, 1980). The oldest belt consists of late Middle–Late Jurassic plutons in the Wallowa terrane (Unruh et al., 2008) and Baker terrane (Fig. 1) (Parker et al., 2008; J. Schwartz, unpublished data). These plutons are typically <3 km<sup>2</sup> in areal extent and range in composition from gabbro to quartz diorite. Geochemically, they are magnesian, calcic to calc-alkalic, and metaluminous, and characterized by low Na, Al, Sr, but high Y concentrations (Fig. 4). This phase of magmatism is coeval with a short-lived episode of regional contraction in the Baker terrane that is bracketed between 159 and 154 Ma (D<sub>2</sub> of Avé Lallemant, 1995; Schwartz and Snoke, 2008).

A second distinct pulse of magmatism occurred between ca. 148 and 137 Ma and consists of two spatially and geochemically distinct belts of high Sr/Y (high Na, Al, Sr, but low Y) plutons in the Greenhorn subterrane of the Baker terrane, and low Sr/Y (low Na, Al, Sr, but high Y) plutons and batholiths in the Bourne subterrane of the Baker terrane and the Wallowa terrane. In the




Figure 3. Histogram of U-Pb zircon ages showing the distribution of Middle Jurassic to Early Cretaceous magmatic rocks in the Blue Mountains Province. Data compiled from Walker (1986; 1989), Manduca (1993), Lee (2004), McClelland and Oldow (2007), Parker et al. (2008), Unruh et al. (2008), and J. Schwartz and K. Johnson (unpublished data).

TABLE 1. SUMMARY OF CHARACTERISTICS OF MIDDLE JURASSIC–EARLY CRETACEOUS PLUTONIC BELTS IN THE BLUE MOUNTAINS PROVINCE, NORTHEAST OREGON

Setting (terrane/subterrane)	Rock type	Age (Ma)	Associated volcanics?	Associated mafic rocks?	Fe-number	MAFI	Na <sub>2</sub> O/K <sub>2</sub> O	~Sr	Sr/Y (ave.)	La/Lu (ave.)	Residual minerals
LoSY Wallowa and Baker terrane	hbl gabbro, hbl ± biotite diorite	162–154	Unknown	Yes	bimodal: ferroan/magnesian	bimodal: alkalic/calcic	4.4	300	12	33	pyroxene, plagioclase
LoSY Wallowa and Baker terrane (Bourne subterrane)	gabbro, hbl diorite, tonalite, granodiorite, granite	148–137	No	Yes	magnesian	calcic-calc alkalic	3.1	469	39	61	pyroxene, plagioclase
HiSY Baker terrane (Greenhorn subterrane)	biotite ± muscovite tonalite-granodiorite	147–145	No	No	magnesian	calcic-calc alkalic	2.7	612	57	120	Amphibole, garnet (±plagioclase)

Note: LoSY—low Sr/Y; HiSY—high Sr/Y; hbl—hornblende; MAFI—modified alkali–lime index.

Bourne subterrane and Wallowa terrane, low Sr/Y plutons and batholiths (148–137 Ma) intrude island-arc and related metasedimentary rocks and are similar in composition to older Middle–Late Jurassic plutons (162–154 Ma belt). In contrast, high Sr/Y plutons in the Greenhorn subterrane (147–145 Ma) spatially overlap the older low Sr/Y plutons, but are much more compositionally restricted (Fig. 4; Table 1).

The transition from low Sr/Y (162–154 Ma belt) to high Sr/Y magmatism (147–145 Ma) in the Greenhorn subterrane occurred ~7 Ma after regional contraction, and raises fundamental questions regarding the origin of high Sr/Y plutons in the Baker terrane. Several geochemical characteristics of the high Sr/Y magmas suggest an origin for these rocks by partial melting in the presence of a plagioclase-poor to absent, hornblende + garnet-bearing source (depths >40 km). These features include: steeply fractionated rare earth element (REE) patterns, high Sr/Y values, and little to no Eu anomaly. Does the observed change in geochemistry from low Sr/Y to high Sr/Y reflect partial melting of orogenically thickened crust as a direct result of Late Jurassic orogenesis?

The last stage of magmatism in the region (pre-Idaho batholith emplacement) consists of small plutons of tonalitic and trondhjemitic composition. This group can be subdivided into two sub-groups: (1) 124–120 Ma metaluminous hornblende-biotite tonalite plutons in a northeast-southwest-trending belt located east (inboard) of the Late Jurassic belt, and (2) 125–111 Ma, strongly peraluminous tonalite and trondhjemite plutons that mostly occur in a belt subparallel to the initial Sr isotopic 0.706 line in western Idaho. These plutons were emplaced >10 Ma after Late Jurassic–Early Cretaceous magmatism in the Baker terrane (146–137 Ma) and represent a distinct phase of magmatism that postdates regional contraction (e.g., Snee et al., 1995; Gray and Oldow, 2005) and peak metamorphism (550–600 °C, 8–9 kbar) at  $128 \pm 3$  Ma in the Salmon River suture zone (Selverstone et al., 1992; Getty et al., 1993).

## REGIONAL TECTONIC IMPLICATIONS

Rocks examined during this field trip highlight an apparent temporal partitioning of sedimentation and magmatism, and




Figure 4. Representative geochemical data for high Sr/Y (HiSY) and low Sr/Y (LoSY) plutons. (A) Both suites are magnesian; however, high Sr/Y plutons have more restricted SiO<sub>2</sub> compositions. (B) Both suites are calcic to calc-alkalic. (C) High Sr/Y plutons have strongly fractionated REE patterns. (D) High Sr/Y plutons have high Sr/Y values (>40) and low Y values (<20 ppm), whereas low Sr/Y plutons have low Sr/Y values (<40) and higher Y values (15–60 ppm). Data from J. Schwartz and K. Johnson (unpublished data).



suggest a systematic shift in the locus of crustal thickening, sedimentation and magmatism through time. The majority of Mesozoic sedimentation appears to have taken place from Late Triassic (ca. 225 Ma) to late Middle Jurassic time (ca. 160 Ma; Dickinson and Thayer, 1978; LaMaskin, 2009). In contrast, magmatism in the Baker terrane appears to have initiated (?) in late Middle Jurassic time (ca. 162 Ma) and continued into the Cretaceous.

We propose a model in which Jurassic sedimentation in the Blue Mountains occurred in a subsiding collisional basin adjacent to an area of crustal thickening in western Nevada (i.e., the Luning-Fencemaker fold-and-thrust belt; Wyld and Wright, 2001; Dorsey and LaMaskin, 2007, 2008). Sedimentation in the Blue Mountains ceased when crustal thickening stepped outboard to a location in the Blue Mountains (Greenhorn subterrane of the Baker terrane; Ferns and Brooks, 1995). This shift in the locus of crustal thickening appears to be recorded by a regional contraction event at ca. 159–154 Ma (i.e.,  $D_2$  of Avé Lallemant, 1995; Schwartz and Snoke, 2008) and by the emplacement of high Sr/Y plutons in Late Jurassic to Early Cretaceous time. Subsequently, the locus of crustal thickening in the region shifted back to the east (i.e., the Salmon River suture zone) and was closely followed by intrusion of small Early Cretaceous tonalitic and trondhjemitic plutons.

## DAY 1

### Directions to Stop 1-1

Begin field trip directions at the intersection of U.S. Highway 26 and U.S. Highway 395 in John Day, Oregon (Fig. 2A). Travel south on U.S. Highway 395 ~18 miles to the intersection with County Route 63 (Izee-Paulina Lane). Turn right on County Route 63 and travel ~18.4 miles, then turn right onto Forest Service Road (FSR) 6370. Continue on winding roads of FSR 6370 for ~15 miles. Turn right onto FSR 304, a small jeep trail located at UTM coordinates 307072E, 4891572N (UTM zone 11, datum NAD 83), and follow ~0.3 miles. Park at the entrance to a closed FSR at UTM coordinates 307022E, 4891985N. On foot, follow closed FSR due west ~1.2 miles, bearing left at intersection with FSR 328, to unvegetated hills of serpentinite on the north side of the road (end hike on road at UTM coordinates 305460W, 4892121N). Hike ~0.25 mi due north to chromite mine pits at UTM coordinates 305393W, 4892252N.

### Stop 1-1. Upper Triassic Vester Formation and Rocks of the Baker Terrane

Throughout the Izee region of central Oregon, the Baker terrane forms the basement to the Triassic–Jurassic basinal assemblage. Upper Triassic sedimentary rocks of the Vester Formation (megasequence 1) are the oldest deposits of the Triassic basinal assemblage, and this is a key locality where they rest in depositional contact on the Baker terrane (Fig. 2). At this location we will observe field evidence for the depositional contact between

the Baker terrane and overlying Vester Formation, and view typical rock types of the Vester Formation (participants are referred to Brown and Thayer, 1977: U.S. Geological Survey [USGS] Map I-1021).

In this area, rocks of the Baker terrane include two dominant rock types: (1) strongly sheared serpentinite, and (2) metavolcanic greenstone (Brown and Thayer, 1977). In the immediate vicinity of Stop 1-1, serpentinite has been mined for chromite and abandoned mine pits can be found at several locations. The Vester Formation has been divided into two members: (1) the Begg Member (~2500 m) is dominantly chert-rich conglomerate and sandstone with lesser amounts of black shale, polymict breccia, and local volcanic rocks, and (2) the Brisbois Member (~1250 m) is dominantly thin-bedded chert-grain sandstone with local calcarenite and limestone olisostromes (Dickinson and Vigrass, 1965; Brown and Thayer, 1977; Dickinson and Thayer, 1978). Paleocurrent indicators reported by Dickinson and Thayer (1978) suggest southeast-directed transport (modern coordinates). The age of the Begg Member is poorly constrained and may be Late Triassic (Carnian) or older, whereas diagnostic pelycepod and ammonite faunas in the Brisbois Member suggest a Late Triassic, Carnian age (*subbullatus* zone; Dickinson and Vigrass, 1965; Blome, 1984; Blome et al., 1986).

The sandstone detrital mode of the Vester Formation is noteworthy for abundant chert and meta-chert grains in conjunction with a relatively low abundance of monocrystalline quartz grains (Dickinson et al., 1979). The unique framework-grain assemblage, paleocurrent indicators suggesting transport away from likely source rocks in the Baker terrane, and numerous syn-depositional faults, indicate that the Vester Formation was derived from uplifted rocks of the Baker terrane during Late Triassic deformation and related basin formation (Dickinson and Thayer, 1978; Brown and Thayer, 1977; Dickinson, 1979; Avé Lallemant, 1995).

Mapping by Brown and Thayer (1977) at this location shows that the Begg Member of the Vester Formation rests in depositional contact on both serpentinite and metavolcanic greenstone of the Baker terrane. The contact is highly irregular and nonplanar, unlike a fault contact, and the mapped attitudes of bedding within the Vester Formation generally follow the Vester-Baker contact. Following a brief orientation, participants will spend some time exploring the contact between rocks of the Baker terrane and the Vester Formation and viewing exposures of chert-pebble conglomerate and chert-grain sandstone.

Medium-grained, chert-rich sandstone samples from the Begg and Brisbois members of the Vester Formation yield abundant detrital-zircon grains (Fig. 5). The majority are well-rounded pink, yellow, or green grains. The samples include minor quantities of Mesozoic-age grains (~2% of grains) and are dominated by Paleozoic- (~41%) and Precambrian-age grains (57%). Prominent peaks in the probability distribution for Mesozoic and Paleozoic ages are present at 374 Ma (Devonian), 311 Ma (Pennsylvanian), and 274 Ma (Permian). The majority of Precambrian-age grains are late Paleoproterozoic in age and fall



within the range of ca. 2.05–1.7 Ga with peaks present at 1.87 and 1.99 Ga. Late Archean ages are represented by grains in the range of 2.75–2.5 Ga.

### Directions to Stop 1-2

Return on foot to vehicles and backtrack to Forest Service Road 6370. Turn left onto FSR 6370 and travel 0.4 miles to small exposures of limestone on the north side of the road (UTM coordinates 307590E, 4891144N).

### Stop 1-2. Upper Triassic Rail Cabin Argillite at Graylock Butte

Here we will make a brief stop to examine bioclastic limestone and calcareous quartzose sandstones and pebble conglomerates of the Rail Cabin Argillite (Fig. 2A). In this area, Upper Triassic and Lower Jurassic rocks that rest on the Vester Formation consist of a relatively thin (~760 m) succession of shallow-water facies including limestone lenses containing well-preserved specimens of *Halorella* that suggest shoaling of the former deep-marine setting (Brown and Thayer, 1977). Correlative rocks on the east side of the Poison Creek fault approach 10,000 m in thickness. Thus syn-depositional motion on the Poison Creek fault appears to have exerted a major structural control on deposition, with uplift in the west and subsidence in the east during Late Triassic to Early Jurassic time (Brown and Thayer, 1977; Dickinson and Thayer, 1978).

Sandstone and pebble conglomerate samples at this locality contain distinctive quartz grains and clasts that display a wide range of sedimentary and metamorphic fabrics. These include grains of quartz arenite sandstone, recrystallized quartz displaying polygonal grain boundaries and stretched and flattened fabrics (i.e., stretched metamorphic quartz). These types of quartz grains and clasts are also present in sediments of the Aldrich Mountains Group east of the Poison Creek fault (Fields Creek Formation), but they are not observed in the underlying Vester Formation. This suggests erosion of a metamorphic source area, and supports a model for progressive regional orogenesis in the Baker terrane during Upper Triassic and Lower Jurassic time (i.e., Dorsey and LaMaskin, 2007; LaMaskin et al., 2008).




Figure 5. Combined detrital-zircon age distributions from the Begg ( $n = 105$ ) and Brisbois ( $n = 112$ ) members of the Vester Formation, representative of megasequence-1 rocks. Data from LaMaskin (2009).

### Directions to Stop 1-3

Continue traveling east on FSR 6370 4.1 miles to prominent exposures on the north side of the road (UTM coordinates 311540E, 4889821N).

### Stop 1-3. Upper Triassic Fields Creek Formation (Aldrich Mountains Group) Bouldery Mudstone and Olistostromal Slide Blocks

We have crossed over the Poison Creek fault (poorly exposed in this area) and passed into exposures of the Fields Creek Formation, the lower unit of the Aldrich Mountains Group (Fig. 2). Rocks at this stop include an anomalous unit of bouldery mudstone and olistostromal slide blocks of limestone and quartz sandstone-conglomerate. To the north, exposures of the Fields Creek Formation are more typical and include thin- to medium-bedded turbidite sandstones.

The Aldrich Mountains Group rests locally in depositional contact on serpentinite-matrix mélange of the Baker terrane (Fig. 2; Dickinson and Thayer, 1978). The Fields Creek Formation (~4500 m) is a thick turbiditic succession of fine-grained sandstone and black shale with slide breccias and olistostromes of igneous and sedimentary rock types in the lower part (Brown and Thayer, 1977; Dickinson and Thayer, 1978). Radiolarian faunas from the Fields Creek Formation indicate a Late Triassic, Norian age and suggest reworking of older Middle to Late Triassic deposits (Blome et al., 1986; Yeh, 1989). The uppermost unit of the Aldrich Mountains Group is the Keller Creek Shale (~1500 m) which includes volcanoclastic greywacke and is Early Jurassic in age (Brown and Thayer, 1977; Imlay, 1986).

A large olistostromal block of quartzose sandstone and conglomerate is present just past where the road turns to the south (when driving east). The lithology is identical to sandstone and conglomerate in the Rail Cabin Argillite west of the Poison Creek fault (Stop 1-2). Quartzose grains and clasts display a wide range of sedimentary and metamorphic fabrics. Other basal sections of the Fields Creek Formation along Fields Creek road to the north contain slide blocks and olistostromes of plutonic rocks identical to small plutons in the Baker terrane (Carpenter and Walker, 1992; LaMaskin, 2009). These relationships and recycled radiolarian faunas (Blome, 1984; Yeh, 1989) show that the Fields Creek Formation was eroded from both the uplifted Baker terrane and from Triassic sedimentary rocks of the Rail Cabin Argillite, and deposited in a complex fault-bounded basin at the margin of a large active thrust belt (Baker terrane).

### Directions to Stop 1-4

Continue traveling east on FSR 6370 ~3.4 miles to outcrops of gray calcareous sandstone on the north side of the road (UTM coordinates 314780E, 4889265N). Fossil-bearing outcrops are present at ~3.4 to 3.7 miles from Stop 1-3.

**Stop 1-4. Lower Jurassic Suplee Formation—Fossil Collecting Locality**

We have now crossed over the regional unconformity between megasequence 1 and overlying sedimentary rocks of megasequence 2 (Fig. 2). Megasequence-2 deposits record a second major phase of subsidence and sedimentation in the Izee region. Sediments are dominantly volcanoclastic and were derived predominantly from inboard locations to the east and northeast (modern coordinates; Dickinson and Thayer, 1978). The basal portion of megasequence 2 consists of the transgressive, shallow- to deep-marine Mowich Group (~690 m; Brown and Thayer, 1977). We will stop to view a 10–20-m-thick, bioturbated calcareous sandstone unit in the Suplee Formation. The Suplee Formation is mainly Pliensbachian age (Imlay, 1968) and is richly fossiliferous. Numerous fossils have been collected at localities in the Suplee Formation, including ammonites, pelycypods, brachiopods, and gastropods.

Turbidite sandstone of the Hyde Formation overlies the Suplee Formation and contains abundant detrital-zircon grains (Fig. 6A). All zircon grains are Mesozoic and define a unimodal probability distribution that spans 191–172 Ma, yielding an Early Jurassic peak at ca. 180 Ma.

**Directions to Stop 1-5**

Continue traveling east on FSR 6370 back to the intersection with County Route 63. Turn left on Route 63 and travel ~6.1 miles. Turn left onto National Forest Road 24 and travel ~1.0 mile to prominent exposures on the east side of the road (UTM coordinates 321958E, 4894830N).

**Stop 1-5. Middle Jurassic Snowshoe Formation along Wikiup Creek**

The final stop of the day is in Middle Jurassic turbidites of the Snowshoe Formation (Fig. 2), stratigraphically higher in




Figure 6. Detrital-zircon age distributions from megasequence-2 rocks. Curves represent the summation of individual detrital-zircon ages and associated 2-sigma Gaussian errors; associated Y-axes represent relative probability. Histograms are based on individual detrital-zircon grain ages and do not incorporate errors; associated Y-axes represent number of grains in a given age bin. Shaded insets show additional detail. (A) The Hyde Formation, Lower Jurassic rocks at the base of megasequence 2. (B) The Snowshoe Formation, Middle Jurassic rocks within megasequence 2. (C) The Lonesome Formation, Middle to Upper(?) Jurassic rocks at the exposed top of megasequence 2. Data from LaMaskin (2009).

megasequence 2. The Snowshoe Formation is mainly Middle Jurassic in age (Imlay, 1986) and is dominated by volcanoclastic turbidites such as those seen at this locality. Here, interbedded mudstone, siltstone and gray sandstone are characteristic of the South Fork Member of the Snowshoe Formation (Smith, 1980). A weighted mean age of ca. 167 Ma for the youngest group of two or more detrital-zircon grains overlapping at 1-sigma for this locality suggests a depositional age at the Bathonian-Bajocian boundary, consistent with the biostratigraphic age of the South Fork Member (Smith, 1980; Imlay, 1986; LaMaskin, 2009).

Detrital-zircon grains from this locality are both euhedral-prismatic, clear-colorless and sub-rounded, pink to yellow grains. The sample includes ~64% Mesozoic, 5% Paleozoic, and 31% Precambrian ages (Fig. 6B). Mesozoic-age zircons form a complex, multimodal age distribution with peaks and shoulders in relative probability between 207 and 168 Ma. Abundant Mesoproterozoic grains define peaks at ca. 1.46, 1.29, 1.15 and 1.04 Ga. A Paleoproterozoic peak is present at ca. 1.84 Ga. Additional sandstone samples collected from megasequence-2 turbidite sandstone of the Lonesome Formation yield detrital-zircon ages similar to those found in the Snowshoe Formation (Fig. 6C). Precambrian grains are dominated by Neoproterozoic and Mesoproterozoic ages with fewer Paleoproterozoic and only scattered Archean ages. A comparison of Precambrian detrital-zircon ages from the Triassic Vester Formation to the Jurassic Snowshoe and Lonesome formations indicates the clear difference in provenance between outboard-derived sands (Vester Formation) and inboard-derived sands (Fig. 7; Snowshoe and Lonesome formations).

End of Day 1.




Figure 7. Comparison of >200 Ma detrital-zircon age distributions from megasequence-1 and megasequence-2 rocks. Note the dramatic difference between detrital-zircons derived from the outboard Baker terrane (megasequence 1) and from inboard locations (megasequence 2). Data from LaMaskin (2009). Megasequence-2 plot includes unpublished data from the Lonesome Formation provided by J. Wright.

## DAY 2

### Directions to Stop 2-1

From John Day, drive east on E. Main Street/U.S.-26 and continue for 30.8 miles to the northeast. Turn left at National Forest Development Road (NFD) 2600-437. Drive 1.2 miles. Turn right (north) on NFD 369. Drive 0.5 miles and park at intersection with NFD 352. Hike 0.15 miles N30°E to NFD 301 (UTM coordinates 370710E, 4932988N).

### Stop 2-1. Basal Section of the Dixie Butte Meta-Andesite Complex

At this stop, we will examine the basal section of the Dixie Butte Meta-andesite (Fig. 8). The Dixie Butte Meta-andesite is exposed over >80 km<sup>2</sup>, making it one of the largest continuous volcanogenic complexes in the Blue Mountains Province (Brooks et al., 1984; Ferns and Brooks, 1995). It consists of a lower sequence of tuffaceous sedimentary rocks and volcanoclastic breccias, and an upper sequence of flow rocks and subordinate volcanoclastic breccias and sills.

The basal portion of the Dixie Butte Meta-andesite is best exposed on Dad's Creek (this stop) and consists of ~2000–2200 ft of thick-bedded andesitic lithic tuff, lithic-clast volcanoclastic breccia, tuffaceous sandstone, graphitic argillite, and pebble conglomerate. Dominant lithic clasts in the pebble conglomerates are trachytic basaltic andesite with subordinate diorite, microdiorite, reworked lithic tuff, and fine-grained chert and argillite (Fig. 9A). Lithic clasts are subrounded to angular in shape. Volcanogenic sandstones locally display normally graded bedding. Argillites and fine- to medium-grained sandstones grade upsection into coarser, lithic-clast volcanoclastic breccias. These rocks are in turn overlain by volcanic flows and tuffs in the upper portion of Dixie Butte (e.g., Fig. 9B; see Stop 2-4).

The relationship between Dixie Butte Meta-andesite and older units (e.g., serpentinite-matrix mélangé, Badger Creek unit [Upper Triassic–Permian]) is poorly understood. Nearby chert argillites similar to those found at the base of Dixie Butte (this stop) contain Wolfcampian (i.e., Late Permian) radiolarians which provide a maximum age for the Dixie Butte Meta-andesite complex. Middle to Late Jurassic plutons (Stop 2-2) intrude the Dixie Butte Meta-andesite. Together these relationships suggest an age between Late Permian and Middle to Late Jurassic. Ongoing detrital zircon work and targeted U-Pb zircon dating of andesites are aimed at further constraining the age of the Dixie Butte Meta-andesite complex.

### Directions to Stop 2-2

Backtrack to U.S.-26. Turn left and continue 2.1 miles northeast. Turn left at NFD 2610. Drive 150 feet and turn left at NFD 096. Continue 0.9 miles (UTM coordinates 372006E, 4933394N).

### Stop 2-2. Dixie Summit Pluton: 162 Ma, Low Sr/Y Magmatism

At this stop, we will examine the Dixie Summit pluton in the eastern portion of the Dixie Butte area (Fig. 8), where late Middle Jurassic plutonic rocks intrude serpentinite-matrix mélangé and volcanic and related rocks of the Dixie Butte Meta-andesite complex. The Dixie Summit pluton ranges in composition from gabbro to quartz diorite, and contains the following primary igneous mineral phases: clinopyroxene + plagioclase feldspar  $\pm$  orthopyroxene  $\pm$  zircon  $\pm$  apatite. Ophitic texture is common in gabbroic

and dioritic samples (e.g., Fig. 9C). The primary igneous minerals are extensively altered at and/or near dike and dikelet contacts by secondary minerals such as chlorite, calcite, albite, or actinolite.

Geochemically, the Dixie Summit plutonic rocks are meta-luminous, magnesian, and calcic as defined by the classification of Frost et al. (2001) and Frost and Frost (2008). They display high  $\text{Al}_2\text{O}_3$  (>19 wt%), CaO (>10 wt%) and positive Eu anomalies suggestive of plagioclase accumulation. They also display slight light rare earth element (LREE) enrichment ( $\text{La/Lu} < 30$ ) and low Sr/Y (<26) values. A diorite from this location yielded a Pb/U age of  $162.0 \pm 2.9$  Ma (J. Schwartz, unpublished data).




Figure 8. Geologic map of the Dixie Butte area, Greenhorn subterrane of the Baker terrane, showing location of field-trip stops for Day 2. Dixie Butte (meta)andesites and related volcaniclastic rocks are intruded by two geochemically distinct suites of low Sr/Y plutons at ca. 162–157 Ma, and high Sr/Y plutons at ca. 146 Ma. Modified from Brooks et al. (1984).



Several features suggest a possible consanguineous relationship between the Dixie Summit pluton and the Dixie Butte Meta-andesite. Potential comagmatic relationships include: (1) the intrusion of 162 Ma gabbro-diorite of the Dixie Summit pluton into basaltic andesite; (2) fragments of gabbro, diorite, and microdiorite in lithic-clast volcanoclastic breccias throughout the Dixie Butte Meta-andesite; (3) intrusion of fine-grained plagioclase-phyric meta-andesite dikes and dikelets into the Dixie Summit pluton and other similar-age plutons at  $161.6 \pm 2.0$  Ma (J. Schwartz, unpublished data); and (4) complementary positive Eu anomalies in the Dixie Summit and negative Eu anomalies in the Dixie Butte Meta-andesite as well as other major-element and trace-element geochemical characteristics suggestive of a cogenetic relationship (compare with Dixie Butte Meta-andesite geochemical data: Stop 2-4).

Other Middle to Late Jurassic plutons in the Dixie Butte area display similar geochemical characteristics to the Dixie Summit pluton (e.g., small plutons and plugs north of Dixie Creek). However, along Standard Creek, a Late Jurassic pluton dated at  $157.7 \pm 1.5$  Ma (Parker et al., 2008) displays ferroan and alkalic geochemical characteristics, suggestive of at least two distinct sources for Middle-Late Jurassic magmatism and/or different modes of differentiation (open-versus closed-system) in the Dixie Butte area.

### Directions to Stop 2-3

Backtrack to U.S.-26. Turn right and continue 9.0 miles southwest. In Prairie City, turn right at NW Johnson Avenue which becomes Dixie Creek Road. Continue 4.6 miles (UTM coordinates 364462E, 4931445N).




Figure 9. Photomicrographs of metavolcanic and plutonic rocks of the Dixie Butte area. (A) Crystal lithic-clast volcanoclastic breccia contains abundant angular to subrounded trachytic basaltic andesite clasts and detrital augite and plagioclase, with minor chert, graphitic argillite, orthopyroxene, quartz, and carbonate grains. (B) Representative example of the Dixie Butte Meta-andesite flow rock displaying porphyritic texture consisting of large plagioclase phenocrysts in a fine-grained matrix of clinopyroxene and plagioclase microlites. (C) Dixie Summit diorite (ca. 162 Ma) displays ophitic texture with resorbed (?) plagioclase surrounded by clinopyroxene. (D) Dixie Creek biotite-hornblende tonalite (ca. 146 Ma) has seriate texture consisting of large plagioclase feldspar and quartz with minor hornblende and biotite.

### Stop 2-3. Dixie Creek Pluton: 146 Ma High Sr/Y Magmatism

At this stop, we will examine the Dixie Creek pluton, which is characteristic of latest Jurassic (ca. 146 Ma) plutonism in the Greenhorn subterrane of the Baker terrane (see 146 Ma plutons in Fig. 3). The Dixie Creek pluton intrudes meta-volcanic rocks of the Dixie Butte Meta-andesite and Badger Creek metasedimentary unit (Ferns and Brooks, 1995), which contains Permian–Late Triassic fauna (Brooks et al. 1984; Ferns and Brooks, 1995). The Dixie Creek pluton consists of biotite-hornblende tonalite with minor K-feldspar and accessory zircon, sphene, and apatite (e.g., Fig. 9D). Graphic intergrowths of quartz and plagioclase are common in some rocks. Secondary (deuteric?) alteration of biotite to chlorite is typical, but minor, in samples of this plutonic unit.

In contrast to the 162–157 Ma plutons in the Dixie Butte area, the Dixie Creek pluton is more compositionally restricted (Figs. 4A and 4B), and much less affected by low-temperature (greenschist-facies) metamorphism. Dixie Creek plutonic rocks displays steeply fractionated REE patterns (Fig. 4C), lack Eu anomalies, and have elevated Sr concentrations and Sr/Y values (typically >600 ppm and >40, respectively; Fig. 4D). These features are fundamentally distinct from the older, low Sr/Y 162–157 Ma suite, and represent a change in the character of magmatism following regional contraction at 159–154 Ma (Schwartz and Snoke, 2008).

#### Directions to Stop 2-4

Drive 0.8 miles north along Dixie Creek Road until road forks. Park (UTM coordinates 365140E, 4932498).

### Stop 2-4. Volcanic Rocks of the Dixie Butte Meta-Andesite Complex

At this stop, we will examine representative volcanic rocks of the Dixie Butte Meta-andesite. This outcrop consists of trachytic plagioclase-phyric basaltic andesite, and is characteristic of the upper section of the Dixie Butte Meta-andesite. Locally, the upper section consists of green to gray, plagioclase- and augite-phyric basaltic andesite and andesite flows, subordinate volcanoclastic breccias, keratophyre, dark-gray basalt, and pale-green silicic flows and tuffs (Brooks et al., 1984; Ferns and Brooks, 1995). The primary igneous mineral phases of the volcanic rocks are plagioclase and augite  $\pm$  orthopyroxene  $\pm$  quartz (e.g., Fig. 9B). Vesicles are common in some rocks and are typically filled with calcite, quartz, and/or chlorite. No pillow structures have been observed. Alteration is typically less intense than other areas of the Greenhorn subterrane (e.g., Ferns and Brooks, 1995) and is highly localized at the meter scale. Low-grade metamorphism of these rocks may be related to hydrothermal fluid flow associated with Jurassic and/or Tertiary magmatism in the area (Mark Ferns, 2008, personal commun.). Alteration miner-

als include: chlorite + albite + calcite + actinolite. Flow rocks and volcanoclastic breccias in the upper section are intruded by fine-grained microgabbro and microdiorite sills and dikes with chilled margins. Andesitic and silicic dikes locally crosscut flows, sills, and tuffaceous rocks, as well as the late Middle Jurassic (ca. 162 Ma) plutonic rocks (see Stop 2-2). One such dike was dated at ca. 162 Ma (see Stop 2-2). No dikes have been observed cross-cutting the 146-Ma high Sr/Y plutons.

Geochemically, volcanic rocks of the Dixie Butte Meta-andesite range from ~53 to 56 wt% SiO<sub>2</sub> and are characterized by high Al<sub>2</sub>O<sub>3</sub> (16–22 wt%), CaO (6–10 wt%) and Y (16–27 ppm), Ni (70–190 ppm), and Cr (150–480 ppm) values, moderate TiO<sub>2</sub> (1.0–1.5 wt%) and MgO (3–9 wt%) values, and low Sr (<365 ppm), Nb (<9 ppm), Sr/Y (<16) and La/Lu (<40) values. Overall, these geochemical features are similar to late Middle Jurassic (162 Ma) plutons in the Dixie Butte area (e.g., Stop 2-2).

#### ACKNOWLEDGMENTS

Funding for our research in the Blue Mountains Province has been provided by the National Science Foundation to Dorsey (EAR-0537691), Vervoort (EAR-0537913), and Snoke and C.D. Frost (EAR-0711470). LaMaskin acknowledges grants from the Geological Society of America, Sigma Xi, and University of Oregon. Jim Wright (University of Georgia) provided additional data from the Lonesome Formation for use in Figure 7. This manuscript benefited from constructive and insightful reviews by Cal Barnes, Mark Ferns, and the GSA editorial staff.

#### REFERENCES CITED

- Armstrong, R.L., Taubeneck, W.H., and Hales, P.O., 1977, Rb-Sr and K-Ar geochronometry of Mesozoic granitic rocks and their Sr isotopic composition, Oregon, Washington, and Idaho: Geological Society of America Bulletin, v. 88, p. 397–411, doi: 10.1130/0016-7606(1977)88<397:RAKGOM>2.0.CO;2.
- Ashley, R.P., 1995, Petrology and deformation history of the Burnt River Schist and associated plutonic rocks in the Burnt River Canyon area, northeastern Oregon, in Vallier, T.L., and Brooks, H.C., eds., Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region: U.S. Geological Survey Professional Paper 1438, p. 457–496.
- Avé Lallemant, H.G., 1995, Pre-Cretaceous tectonic evolution of the Blue Mountains Province, northeastern Oregon, in Vallier, T.L., and Brooks, H.C., eds., Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region: U.S. Geological Survey Professional Paper 1438, p. 271–304.
- Blome, C.D., 1984, Upper Triassic Radiolaria and Radiolarian zonation from western North America: Bulletins of American Paleontology, v. 85, p. 1–88.
- Blome, C.D., Jones, D.L., Murchey, B.L., and Liniecki, M., 1986, Geologic implications of radiolarian-bearing Paleozoic and Mesozoic rocks from the Blue Mountains Province, eastern Oregon, in Vallier, T.L., and Brooks, H.C., eds., Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Geologic implications of Paleozoic and Mesozoic paleontology and biostratigraphy, Blue Mountains Province, Oregon and Idaho: U.S. Geological Survey Professional Paper 1435, p. 79–101.
- Brooks, H.C., 1979, Geologic map of Huntington and part of Olds Ferry quadrangles, Oregon: State of Oregon, Oregon Department of Geology and Mineral Industries, Geologic Map Series, GMS-13, scale 1:62,500.
- Brooks, H.C., and Vallier, T.L., 1978, Mesozoic rocks and tectonic evolution of eastern Oregon and western Idaho, in Howell, D.G., and McDougall,



- K.A., eds., Mesozoic paleogeography of the western United States: Society of Economic Paleontologists and Mineralogists, Pacific Section, Paleogeography Symposium 2, p. 133–145.
- Brooks, H.C., McIntyre, J.R., and Walker, G.W., 1976, Geology of the Oregon part of the Baker 1° × 2° quadrangle: State of Oregon, Oregon Department of Geology and Mineral Industries, Geologic Map Series, GMS-7, scale 1:250,000.
- Brooks, H.C., Ferns, M.L., and Avery, D.G., 1984, Geology and gold deposit map of the southwest quarter of the Bates Quadrangle, Grant County, Oregon: Oregon Department of Geology and Mineral Industries, GMS-35, scale 1:24,000.
- Brown, C.E., and Thayer, T.P., 1966, Geologic map of the Canyon City quadrangle, northeastern Oregon: U.S. Geological Survey, Map I-447, scale 1:62,500.
- Brown, C.E., and Thayer, T.P., 1977, Geologic map of pre-Tertiary rocks in the eastern Aldrich Mountains and adjacent areas to the south, Grant County, Oregon: U.S. Geological Survey, Miscellaneous Investigations Series I-1021, scale 1:62,500.
- Carpenter, P.S., and Walker, N.W., 1992, Origin and tectonic significance of the Aldrich Mountains serpentinite matrix mélange, northeastern Oregon: *Tectonics*, v. 11, p. 690–708, doi: 10.1029/91TC03013.
- Dickinson, W.R., 1979, Mesozoic forearc basin in central Oregon: *Geology*, v. 7, p. 166–170, doi: 10.1130/0091-7613(1979)7<166:MFBCO>2.0.CO;2.
- Dickinson, W.R., 2004, Evolution of the North American Cordillera: *Annual Review of Earth and Planetary Sciences*, v. 32, p. 13–45, doi: 10.1146/annurev.earth.32.101802.120257.
- Dickinson, W.R., and Gehrels, G.E., 2003, U-Pb ages of detrital zircons from Permian and Jurassic eolian sandstones of the Colorado Plateau, USA: Paleogeographic implications: *Sedimentary Geology*, v. 163, p. 29–66, doi: 10.1016/S0037-0738(03)00158-1.
- Dickinson, W.R., and Gehrels, G.E., 2009, U-Pb ages of detrital zircons in relation to paleogeography: Triassic paleodrainage networks and sediment dispersal across southwest Laurentia: *Journal of Sedimentary Research*, v. 78, p. 745–764, doi: 10.2110/jsr.2008.088.
- Dickinson, W.R., and Gehrels, G.E., 2009, U-Pb ages of detrital zircons in Jurassic eolian and associated sandstones of the Colorado Plateau: Evidence for transcontinental dispersal and intraregional recycling of sediment: *Geological Society of America Bulletin*, v. 121, p. 408–433, doi: 10.1130/B26406.1.
- Dickinson, W.R., and Thayer, T.P., 1978, Paleogeographic and paleotectonic implications of Mesozoic stratigraphy and structure in the John Day inlier of central Oregon, *in* Howell, D.G., and McDougall, K.A., eds., Mesozoic paleogeography of the western United States: Society of Economic Paleontologists and Mineralogists, Pacific Section, Paleogeography Symposium 2, p. 147–161.
- Dickinson, W.R., and Vigrass, L.W., 1965, Geology of the Suplee-Izee area, Crook, Grant, and Harney counties, Oregon: Oregon Department of Geology and Mineral Industries Bulletin 58, 109 p.
- Dickinson, W.R., Helmold, K.P., and Stein, J.A., 1979, Mesozoic lithic sandstones in central Oregon: *Journal of Sedimentary Petrology*, v. 49, p. 501–516.
- Dorsey, R.J., and LaMaskin, T.A., 2007, Stratigraphic record of Triassic–Jurassic collisional tectonics in the Blue Mountains Province, northeastern Oregon: *American Journal of Science*, v. 307, p. 1167–1193, doi: 10.2475/10.2007.03.
- Dorsey, R.J., and LaMaskin, T.A., 2008, Mesozoic collision and accretion of oceanic terranes in the Blue Mountains Province of northeastern Oregon: New insights from the stratigraphic record, *in* Spencer, J.E., and Titley, S.R., eds., *Circum-Pacific Tectonics, Geologic Evolution, and Ore Deposits: Tucson, Arizona*, Arizona Geological Society, Digest 22, p. 325–332.
- Ferns, M.L., and Brooks, H.C., 1995, The Bourne and Greenhorn subterranean of the Baker terrane, northeastern Oregon: Implications for the evolution of the Blue Mountains island-arc system, *in* Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region*: U.S. Geological Survey Professional Paper 1438, p. 331–358.
- Ferns, M.L., Madin, I.P., and Taubeneck, W.H., 2001, Reconnaissance geologic map of the La Grande 30' × 60' quadrangle, Baker, Grant, Umatilla, and Union Counties, Oregon: Oregon Department of Geology and Mineral Industries Reconnaissance Map Series RMS-1, map scale 1:24,000, 52 p.
- Frost, B.R., and Frost, C.D., 2008, A geochemical classification for feldspathic igneous rocks: *Journal of Petrology*, v. 49, p. 1955–1969, doi: 10.1093/ptrology/egn054.
- Frost, B.R., Barnes, C.G., Collins, W.J., Arculus, R.J., Ellis, D.J., and Frost, C.D., 2001, A geochemical classification for granitic rocks: *Journal of Petrology*, v. 42, p. 2033–2048, doi: 10.1093/ptrology/42.11.2033.
- Getty, S.R., Selverstone, J., Wernicke, B.P., Jacobsen, S.B., and Aliberti, E., 1993, Sm–Nd dating of multiple garnet growth events in an arc-continent collision zone, northwestern U.S. Cordillera: *Contributions to Mineralogy and Petrology*, v. 115, p. 45–57, doi: 10.1007/BF00712977.
- Gray, K.D., and Oldow, J.S., 2005, Contrasting structural histories of the Salmon River belt and Wallowa terrane: Implications for terrane accretion in northeastern Oregon and west-central Idaho: *Geological Society of America Bulletin*, v. 117, p. 687–706, doi: 10.1130/B25411.1.
- Hamilton, W.B., 1963, Metamorphism in the Riggins region, western Idaho: U.S. Geological Survey Professional Paper 436, 95 p.
- Harding, J.P., Gehrels, G.E., Harwood, D.S., and Girty, G.H., 2000, Detrital zircon geochronology of the Shoo Fly Complex, northern Sierra terrane, northeastern California *in* Soreghan, M.J., and Gehrels, G.E., eds., *Paleozoic and Triassic paleogeography and tectonics of western Nevada and northern California*: Geological Society of America Special Paper 347, p. 43–55, doi: 10.1130/0-8137-2347-7.43.
- Hendricksen, T.A., Skurla, S.J., and Field, C.W., 1972, K–Ar dates for plutons from the Iron Mountain and Sturgill Peak areas of western Idaho: *Isochron-West*, v. 5, p. 13–16.
- Hotz, P.E., Lanphere, M.A., and Swanson, D.A., 1977, Triassic blueschist from northern California and north-central Oregon: *Geology*, v. 5, p. 659–663, doi: 10.1130/0091-7613(1977)5<659:TBFNCA>2.0.CO;2.
- Imlay, R.W., 1986, Jurassic ammonites and biostratigraphy of eastern Oregon and western Idaho, *in* Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Geologic implications of Paleozoic and Mesozoic paleontology and biostratigraphy, Blue Mountains Province, Oregon and Idaho*: U.S. Geological Survey Professional Paper 1435, p. 53–57.
- Johnson, K., and Barnes, C.G., 2002, Jurassic magmatism prior to and during terrane accretion in the Blue Mountains province, NE Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 34, no. 5, p. 21–22.
- Johnson, K., Walton, C., Barnes, C.G., and Kistler, R.W., 1995, Time-dependent geochemical variations of Jurassic and Cretaceous plutons in the Blue Mountains, northeastern Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 27, no. 6, p. 435.
- Jones, D.L., Pessagno, E.A., Jr., Force, E.R., and Irwin, W.P., 1976, Jurassic radiolarian chert from near John Day, Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 8, no. 3, p. 386.
- Kays, M.A., Ferns, M.L., and Brooks, H.C., 1987, Metamorphism of Triassic–Paleozoic belt rocks—A guide to field and petrologic relations in the oceanic mélange, Klamath and Blue Mountains, California and Oregon: *in* Ernst, W.G., ed., *Metamorphism and crustal evolution of the Western United States (Rubey Volume VII)*: Englewood, Cliffs, New Jersey, Prentice-Hall, p. 1098–1120.
- Kays, M.A., Stimac, J.P., and Goebel, P.M., 2006, Permian–Jurassic growth and amalgamation of the Wallowa composite terrane, northeastern Oregon, *in* Snoke, A.W., and Barnes, C.G., eds., *Geological studies in the Klamath Mountains province, California and Oregon: A volume in honor of William P. Irwin*: Geological Society of America Special Paper 410, p. 465–494.
- LaMaskin, T.A., 2008, Late Triassic (Carnian–Norian) mixed volcanoclastic-carbonate facies of the Olds Ferry terrane, eastern Oregon and western Idaho, *in* Blodgett, R., and Stanley, G., eds., *The terrane puzzle: New perspectives on paleontology and stratigraphy from the North American Cordillera*: Geological Society of America, Special Paper 442, p. 251–267.
- LaMaskin, T.A., 2009, Stratigraphy, Provenance, and Tectonic Evolution of Mesozoic Basins in the Blue Mountains Province, eastern Oregon and western Idaho [Ph.D. dissertation]: Eugene, Oregon, University of Oregon, 232 p.
- LaMaskin, T.A., Dorsey, R.J., and Vervoort, J.V., 2008, Tectonic controls on mudrock geochemistry, Mesozoic rocks of eastern Oregon and western Idaho, U.S.A.: Implications for Cordilleran tectonics: *Journal of Sedimentary Research*, v. 78, p. 765–783, doi: 10.2110/jsr.2008.087.
- Lee, R.G., 2004, The geochemistry, stable isotopic composition, and U–Pb geochronology of tonalite-trondhjemites within the accreted terrane, near Greer, north-central Idaho [M.S. thesis]: Pullman, Washington, Washington State University, 132 p.
- Leeman, W.P., Avé Lallemant, H.G., Gerlach, D.C., Sutter, J.F., and Arculus, R.J., 1995, Petrology of the Canyon Mountain Complex, *in* Vallier, T.L.,

- and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and Tectonic Evolution of pre-Tertiary Rocks of the Blue Mountains Region*: U.S. Geological Survey, Professional Paper 1438, p. 1–43.
- Lewis, R.S., 2002, *Geologic maps of Nez Perce, Lewis, Idaho, Adams and Washington counties*: Digital Atlas of Idaho: available from <http://imnh.isu.edu/digitalatlas/> (accessed 2004).
- Lund, K., 2004, *Geology of the Payette National Forest and vicinity, west-central Idaho*: U.S. Geological Survey Professional Paper 1666-A, -B, 89 p.
- Manduca, C.A., Kuntz, M.A., and Silver, L.T., 1993, Emplacement and deformation history of the western margin of the Idaho batholith near McCall: Influence of a major terrane boundary: *Geological Society of America Bulletin*, v. 105 p. 749–765, doi: 10.1130/0016-7606(1993)105<0749:EADHOT>2.3.CO;2.
- Mann, G.M., and Vallier, T.L., 2007, Mesozoic telescoping of island arc terranes and geologic evolution of the Cuddy Mountains region, western Idaho, in Kuntz, M.A., and Snee, L.W., eds., *Geological Studies of the Salmon River Suture Zone and Adjoining Areas, West-Central Idaho and Eastern Oregon*: U.S. Geological Survey, Professional Paper 1738, p. 123–162.
- McClelland, W.C., and Oldow, J.S., 2007, Late Cretaceous truncation of the western Idaho shear zone in the central North American Cordillera: *Geology*, v. 35, p. 723–726, doi: 10.1130/G23623A.1.
- Parker, K.O., Schwartz, J.J., and Johnson, K., 2008, Petrology and age of Dixie Butte plutonic rocks, Blue Mountains, NE Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 40, no. 6, p. 154–155.
- Pessagno, E.A., and Blome, C.D., 1986, Faunal affinities and tectonogenesis of Mesozoic rocks in the Blue Mountains province of eastern Oregon and western Idaho, in Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Geologic implications of Paleozoic and Mesozoic paleontology and biostratigraphy, Blue Mountains province, Oregon and Idaho*: U.S. Geological Survey Professional Paper 1435, p. 65–78.
- Rahl, J.M., Reiners, P.W., Campbell, I.H., Nicolescu, S., and Allen, C.M., 2003, Combined single-grain (U-Th)/He and U/Pb dating of detrital zircons from the Navajo Sandstone, Utah: *Geology*, v. 31, p. 761–764, doi: 10.1130/G19653.1.
- Selverstone, J., Wernicke, B., and Aliberti, E.A., 1992, Intracontinental subduction and hinged unroofing along the Salmon River suture zone, west central Idaho: *Tectonics*, v. 11, p. 124–144, doi: 10.1029/91TC02418.
- Scherer, H.H., 2006, *Field, geochronologic, and geochemical constraints on the early Mesozoic paleogeographic and tectonic evolution of the central Klamath Mountains, California* [Ph.D. dissertation]: Palo Alto, California, Stanford University, 199 p.
- Schwartz, J.J., and Snoke, A.W., 2008, Timing of tectonic juxtaposition of the Bourne and Greenhorn subterrains, Baker terrane, northeast Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 40, no. 6, p. 187.
- Schwartz, J.J., Snoke, A.W., and Frost, C.D., 2006, Isotopic characteristics of the Baker terrane, NE Oregon: Implications for the relationship between the Bourne and Greenhorn subterrains [abs.]: *Geological Society of America Abstracts with Programs*, v. 38, no. 7, p. 449.
- Schwartz, J.J., Snoke, A.W., Frost, C.D., Barnes, C.G., Gromet, L.P., and Johnson, K., 2010, Analysis of the Wallowa-Baker terrane boundary: Implications for tectonic accretion in the Blue Mountains province, northeastern Oregon: *Geological Society of America Bulletin*, doi: 10.1130/B26493.1 (in press).
- Silberling, N.J., Jones, D.L., Blake, M.C., Jr., and Howell, D.G., 1984, Litho-tectonic terrane map of the western conterminous United States: U.S. Geological Survey Open-File Report 84-0523, p. C1–C43.
- Smith, P.L., 1980, Correlation of the members of the Jurassic Snowshoe Formation in the Izee Basin of east-central Oregon: *Canadian Journal of Earth Sciences*, v. 17, p. 1603–1608.
- Snee, L.W., Lund, K., Sutter, J.F., Balcer, D.E., and Evans, K.V., 1995, An  $^{40}\text{Ar}/^{39}\text{Ar}$  chronicle of the tectonic development of the Salmon River suture zone, western Idaho, in Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region*: U.S. Geological Survey Professional Paper 1438, p. 359–414.
- Thayer, T.P., and Brown, C.E., 1966, *Geologic map of the Aldrich Mountain quadrangle, Grant County, Oregon*: U.S. Geological Survey Geological Quadrangle Map GQ-438, scale 1:62,500.
- Tumpene, K., Crowley, J., Schmitz, M., and Northrup, C.J., 2008, New geochronological constraints on the deposition of the Huntington Formation, Olds Ferry terrane, and implications for the evolution of the Blue Mountains Province [abs.]: *Geological Society of America Abstracts with Programs*, v. 40, no. 1, p. 47.
- Unruh, D.M., Lund, K., Kuntz, M.A., and Snee, L.W., 2008, Uranium-lead zircon ages and Sr, Nd, and Pb isotope geochemistry of selected plutonic rocks from western Idaho: U.S. Geological Survey Open-File Report 2008-1142, 37 p.
- Vallier, T.L., 1977, The Permian and Triassic Seven Devils Group, western Idaho and northeast Oregon: U.S. Geological Survey Bulletin 1437, 58 p.
- Vallier, T.L., 1995, Petrology of pre-Tertiary igneous rocks in the Blue Mountains region of Oregon, Idaho, and Washington: implications for the geologic evolution of a complex island arc, in Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region*: U.S. Geological Survey Professional Paper 1438, p. 125–209.
- Vallier, T.L., 1998, *Islands and Rapids: A geologic story of Hells Canyon*: Lewiston, Idaho, Confluence Press, Lewis-Clark State College, 151 p.
- Walker, G.W., and MacLeod, N.S., 1991, *Geologic map of Oregon*: U.S. Geological Survey, scale 1:500,000, 2 sheets.
- Walker, N.W., 1986, *U/Pb geochronologic and petrologic studies in the Blue Mountains terrane, northeastern Oregon and westernmost-central Idaho: Implications for pre-Tertiary tectonic evolution* [Ph.D. dissertation]: Santa Barbara, University of California, 224 p.
- Walker, N.W., 1989, Early Cretaceous initiation of post-tectonic plutonism and the age of the Connor Creek fault, northeastern Oregon [abs.]: *Geological Society of America Abstracts with Programs*, v. 21, no. 5, p. 155.
- Walker, N.W., 1995, Tectonic implications of U-Pb zircon ages of the Canyon Mountain complex, Sparta complex, and related metaplutonic rocks of the Baker terrane, northeastern Oregon, in Vallier, T.L., and Brooks, H.C., eds., *Geology of the Blue Mountains region of Oregon, Idaho, and Washington: Petrology and tectonic evolution of pre-Tertiary rocks of the Blue Mountains region*: U.S. Geological Survey Professional Paper 1438, p. 247–269.
- White, J.D.L., White, D.L., Vallier, T.L., Stanley, G.D., Jr., and Ash, S.R., 1992, Middle Jurassic strata link Wallowa, Olds Ferry, and Izee terranes in the accreted Blue Mountains island arc, northeastern Oregon: *Geology*, v. 20, p. 729–732, doi: 10.1130/0091-7613(1992)020<0729:MJSLWO>2.3.CO;2.
- Wilson, D., and Cox, A., 1980, Paleomagnetic evidence for tectonic rotation of Jurassic plutons in Blue Mountains, eastern Oregon: *Journal of Geophysical Research*, v. 85, p. 3681–3689, doi: 10.1029/JB085iB07p03681.
- Wyld, S.J., 2002, Structural evolution of a Mesozoic backarc fold-and-thrust belt in the U.S. Cordillera: new evidence from northern Nevada: *Geological Society of America Bulletin*, v. 114, p. 1452–1468, doi: 10.1130/0016-7606(2002)114<1452:SEOAMB>2.0.CO;2.
- Wyld, S.J., and Wright, J.E., 2001, New evidence for Cretaceous strike-slip faulting in the United States Cordillera and implications for terrane-displacement, deformation patterns, and plutonism: *American Journal of Science*, v. 301, p. 150–181, doi: 10.2475/ajs.301.2.150.
- Wyld, S.J., Quinn, M.J., and Wright, J.E., 1996, Anomalous(?) Early Jurassic deformation in the western United States Cordillera: *Geology*, v. 24, p. 1037–1040, doi: 10.1130/0091-7613(1996)024<1037:AEJDIT>2.3.CO;2.
- Wyld, S.J., Rogers, J.W., and Copeland, P., 2003, Metamorphic evolution of the Luning-Fencemaker fold-thrust belt, Nevada: illite crystallinity, metamorphic petrology, and  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology: *The Journal of Geology*, v. 111, p. 17–38, doi: 10.1086/344663.
- Yeh, K.Y., 1989, *Studies of Radiolaria from Fields Creek Formation, east-central Oregon, U.S.A.*: National Museum of Natural Science, Taiwan, Bulletin no. 1, p. 43–109.